

## Betamethasone Dipropionate

【TBET】 Betamethasone 0.064% Oint. 10g/Tube

ATC Code : D07AC01

中文名： 貝他每松軟膏 «杏輝»

適應症： 皮膚炎、皮膚過敏及其他皮質類固醇具有療效之皮膚疾病。

【TSEP】 Septon® 0.064% Solution 10g/Bot

ATC Code : D07AC01

中文名： 舒膚通膠液 «鹽野義»

適應症： 濕疹、皮膚炎群（進行性指掌角皮症、威達耳氏 Vidal's 苔癬）、乾癬掌蹠膿皰疹、紅皮症。

藥理分類： Corticosteroid, Topical

用法用量： Apply one to several times daily until improvement occurs.

不良反應： 皮膚發紅、微血管擴張、萎縮、易感染、毛囊炎。

注意事項： 1.外用製劑。切勿內服，勿接觸眼睛。勿使用於傷口及感染部位。  
2.請保存於 30°C 以下陰涼避光處。

懷 孕 期： 1.動物實驗有催畸形作用之報告，孕婦或可能懷孕之婦人以不使用本劑為佳。  
2. Use of mild to moderate potency topical corticosteroids is preferred in pregnant females, and the use of large amounts or use for prolonged periods of time should be avoided (*Chi 2016; Chi 2017; Murase 2014*).  
Also avoid areas of high percutaneous absorption (*Chi 2017*). The risk of stretch marks may be increased with use of topical corticosteroids (*Murase 2014*).

授 乳 期： 1. It is not known if systemic absorption following topical administration results in detectable quantities of betamethasone in breast milk.  
2. However, topical corticosteroids are generally considered acceptable for use (*Butler 2014; WHO 2002*).  
3. Do not apply topical corticosteroids to breast until breastfeeding ceases (*Leachman 2006*); hypertension was noted in a breastfed infant when a high potency topical corticosteroid was applied to the nipple (*Butler 2014; Leachman 2006*).