

Silver Sulfadiazine

【TSILV2】 Siliverzine Cream® 10mg/g, 20g/Tube

ATC Code : D06BA01

【TSILV5】 Siliverzine Cream® 10mg/g, 500g/Bot

ATC Code : D06BA01

中文名： 燙膚舒乳膏 «杏輝»

適應症： 二級及三級燙傷之預防感染燙傷及一般性外傷傷口之感染皮膚粘膜之感染。

藥理分類： Antibiotic, Topical

用法用量： Administration:

For topical use only; avoid contact with eyes. Apply with a sterile-gloved hand. Burned area should be covered with cream at all times; reapply to areas where cream has been removed by patient activity. Dressings may be used if necessary. Reapply immediately after hydrotherapy.

Apply 1-2 times daily.

禁忌： 磺醯胺之治療可提高核質性黃疸之可能性，本品不應使用於懷孕期、早產兒或第一個月的新生兒。

不良反應： 皮膚搔癢、發疹、紅腫。

注意事項： 1.對磺胺類過敏、光敏感、肝臟或腎臟功能不佳者，小心使用。
2.新生兒及早產兒不推薦使用。

懷 孕 期： Because of the theoretical increased risk for hyperbilirubinemia and kernicterus, silver sulfadiazine is contraindicated for use near term, on premature infants, or on newborn infants during the first 2 months of life.

授 乳 期： 1. It is not known if silver sulfadiazine is found in breast milk following topical application.
2. Due to the potential for serious adverse reactions in the nursing infant, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of treatment to the mother.

儲 存： 請避光保存。