

Oxymetazoline HCl

【TSIND】Sindecon[®] Nasal Spray 0.05% 0.5mg/mL, 15mL/Bot

ATC Code : R01AA05

中文名：醫鼻易噴鼻液 «生達化學»

適應症：急性鼻炎、過敏性鼻炎、血管舒縮性鼻炎、鼻竇炎、鼻咽炎。

藥理分類：**Adrenergic Agonist Agent; Decongestant.**

用法用量：**Note:** NOT recommended for use in children <6 years of age (especially in infants) due to CNS depression.

Administration:

- For intranasal use only.
- Before using for the first time, prime the pump by firmly depressing the rim several times.
- Keep head upright and insert nozzle into nostril, depress rim firmly, and inhale deeply.

Dosage regimens:

Nasal Congestion:

Adults & Children > 6 years of age---

One spray in each nostril 2~3 time daily.

Maximum of 2 times (2 doses) in a 24-hour period.

不良反應：灼熱感、針刺感、鼻乾燥、頭痛、失眠、眩暈、顫抖、心悸或反彈性充血。

注意事項：1. 不建議6歲以下孩童使用本品。
2. 使用3天後症狀若依然存在，則請教醫師。

懷 孕 期：1. Adverse fetal/neonatal events have been noted in case reports following large doses or extended use of oxymetazoline nasal spray in the first trimester of pregnancy (*Baxi 1985; Holm 1985; Menezes 2016*).
2. Decongestants are not the preferred agents for the treatment of rhinitis during pregnancy.
3. Short-term (<3 days) use of intranasal oxymetazoline may be beneficial to some patients although its safety during pregnancy has not been studied (*Wallace 2008*).

授 乳 期：If a decongestant is needed, nasal preparations may be preferred over systemic agents in nursing women (*Anderson 2000*).