

Pantoprazole

【IPANT】 Pantyl for I.V. Injection 40mg

ATC Code : A02BC02

中文名： 速潰樂 凍晶注射劑 40 毫克 《中國化學》

適應症： 合併二種適當之抗微生物製劑治療與幽門螺旋桿菌相關之消化性潰瘍、胃潰瘍、十二指腸潰瘍、中度及嚴重逆流性食道炎、Zollinger-Ellison-Syndrome。

藥理分類： Proton Pump Inhibitor.

用法用量： **Administration:**

- IV: Flush IV line before and after administration with D5W, NS, or LR. In-line filter not required.
- **2-minute infusion:** The volume of reconstituted solution (4 mg/mL) may be administered IV over at least 2 minutes.
- **15-minute infusion:** Infuse over ~15 minutes at a rate of ~7 mL/minute.
- **Continuous infusion:** May also be administered as a continuous infusion for the prevention of rebleeding with in peptic ulcer bleed (off-label use).

Erosive esophagitis associated with GERD:

IV: 40 mg once daily for 7 to 10 days.

Hypersecretory disorders (including Zollinger-Ellison):

IV:

80 mg Q12 H;

160 to 240 mg daily in divided doses has been used up to 7 days.

不良反應： 腹瀉、便秘、脹氣、頭痛。

交互作用：

- **ERLOTINIB:** ↓ erlotinib exposure
- **METHOTREXATE:** ↑ concentration of methotrexate.
- **CAPECITABINE:** ↓ capecitabine bioavailability.
- **KETOCONAZOLE:** ↓ ketoconazole exposure.
- **WARFARIN:** ↑ INR and prothrombin time.

注意事項： 1. 針劑只建議用於不適用於口服投予之情況。
2. 注射前後應以相容輸注液沖洗過。

注射劑調製與

相關注意事項： 1. 每一小瓶內注入 10ml 注射用生理食鹽水溶解。配製後的溶液可直接注射投予或以 100ml 注射用生理食鹽水或 5% 之葡萄糖注射液混合滴注。Pantoprazole 不可與上述以外之溶劑混合。
2. 配製後的溶液須於 12 小時內使用。使用靜脈注射投予時須超過 2-15 分鐘。

懷 孕 期： Based on available data, PPIs may be used when clinically indicated (*Body 2016; Matok 2012; Pasternak 2010; van der Woude 2014*).

授 乳 期： 1. Pantoprazole is present in breast milk (*Plante 2004*).
2. No adverse events were reported in the mother or infant for the 2 weeks following administration (*Plante 2004*).
3. The decision to breastfeed during therapy should consider the risk of infant exposure, the benefits of breastfeeding to the infant, and benefits of treatment to the mother.