## **Activated Charcoal**

[OCARB] Carbomix® Granules for Suspension 61.50 g/bot ATC Code: A07BA01

中文名: 蓋抹滅顆粒劑 《科懋》

適應症: 緊急治療藥物或化學品中毒之病人。

用法用量: Administration: Extemporaneously, mix powder with sufficient water (e.g., 20-30 g

in at least 240 mL) to form a slurry.

**Dosage regimens:** 

**Poisonings:** 

-Adults & Adolescents  $\ge 13 \text{ yr}$ :

Single dose: 25-100 g ORALLY.

Multiple dose:

50-100 g ORALLY, then 12.5 g every hour, 25 g every 2 hrs, or 50 g every 4 hrs. Continue multiple-dose therapy until patient recovers or major toxicity resolves.

-Pediatric Patients:

Single-Dose or Multiple-Dose Regimens for Treatment of

Poisoning in Children

| Age                          | Single Dose           | Multiple Dose                             |
|------------------------------|-----------------------|---|
| Infants up to 1 yr of age    | 10-25 g or 0.5-1 g/kg |   |
| Children up to 13 yrs of age | 25-50 g or 0.5-1 g/kg | 10-25 g initially, then<br>1-2 g/kg Q2-4h |

## **GI Disorders:**

Adults, 0.6-5 g ORALLY as a single dose or 0.975-3.9 g ORALLY 3 times daily after meals.

不良反應: 嘔吐、腹瀉、便祕、解黑便。

注意事項: 本劑對腐蝕性物質引起之中毒無效。

授 乳 期: 授乳婦使用本劑對被哺乳的嬰兒不會造成任何影響。 \*\*全民健保藥品給付規定: Carbomix®限藥物中毒急救使用。