Danazol

[ODANA] Danol® Cap (Danazol) 200 mg/Cap ATC Code : G03XA01

中文名: 達爾諾膠囊 《衛達化學》

適應症: 子宮內膜異位症、早發春機發動、囊狀慢性乳腺炎、乳腺肥大、乳痛、月經過

多、纖維性囊腫性乳房疾病。

藥理分類: Androgen.

作用機轉: Suppresses pituitary output of FSH and LH, resulting in regression and atrophy of

normal and ectopic endometrial tissue; decreases rate of growth of abnormal breast

tissue.

用法用量:Administration: Orally, administer consistently with regards to food; either always

administer with food or administer on an empty stomach.

Note: Initiate therapy during menstruation or ensure patient is not pregnant while on therapy.

Indications and dosage regimen:

Endometriosis:

-Moderate to severe cases:

initially, 800 mg in 2 divided doses; adjust depending on patient response.

-Mild cases:

initially, 200-400 mg given in 2 divided doses; adjust depending on patient response.

Therapy for endometriosis must continue uninterrupted for 3-6 months but may be extended for 9 months if necessary.

Fibrocystic breast disease:

100-400 mg daily given in 2 divided doses depending on patient response. When treating fibrocystic breast disease, breast pain and tenderness are significantly relieved by the first month and eliminated in 2-3 months; elimination of nodularity usually requires 4-6 months of uninterrupted therapy.

Hereditary angioedema:

- —Initial, 200 mg 2-3 times daily; individualize based on patient's clinical response.
- Maintenance, decrease initial dose by 50% or less at intervals of 1-3 months or longer; if attack occurs, daily dose may be increased by up to 200 mg.

不良反應: 光敏感皮膚疹、顏面潮紅、出汗、男性化作用、陰道出血。

注意事項: 1.服用本藥期間應定期監測肝功能及做血液檢查。

- 2.服藥期間可能月經不規則或無月經。停止服藥後 60-90 天會開始規則月經,若無,應告知醫師。
- 3.服用本藥後會對陽光過敏,應避免在陽光下太久,並作好防曬措施。
- 4.病人應謹慎注意男性化的現象。
- 5.每日劑量超過 400 mg 會有血清酶升高或黃疸等肝功能損害報告。

懷 孕 期: 孕婦勿用本品。因此治療期間懷孕了,則須停藥,若繼續投與胎兒會有男性化 現象,尤其女胎兒會明顯導致外生殖器陰蒂肥大和陰唇融合。

授 乳 期: 哺乳婦勿用本品。