Dicloxacillin Sodium

[ODIC] Ziefmycin® 250mg/Cap ATC Code: J01CF01

中文名: 奇福黴素膠囊 《永信》

適應症: 葡萄球菌、β-溶血性鏈球菌和肺炎雙球菌等細菌所引起的感染症。

藥理分類: Antibiotic, Penicillin

用法用量: Administration: orally taken at least 1 hr before or 2 hrs after meals.

Indications and dosage regimens:

Staphylococcal Infections:

-Mild to Moderate Infections:

Children < 40 kg: 12.5 mg/kg daily given in divided doses every 6 hrs.

Children \geq 40 kg & Adults: 125 mg every 6 hrs.

Children ≥ 1 month of age: AAP recommends 25-50 mg/kg daily in 4 divided doses

-More Severe Infections:

Children < 40 kg:

25 mg/kg daily given in divided doses every 6 hrs; higher dosage may be necessary depending on severity of infection.

Children \geq 40 kg & Adults:

250 mg every 6 hrs; higher dosage may be necessary depending on severity of infection.

-Acute or Chronic Osteomyelitis:

50-100 mg/kg/day divided every 6 hrs have been used.

不良反應: 過敏性反應、間質性腎炎、腹瀉、噁心、腹痛、肝炎。

注意事項: 建議空腹服用(飯前一小時或飯後2小時)。

懷 孕 期: 1.藥物會通過胎盤。安全性尚未建立,用於孕婦需謹慎評估。

2. Adverse events have not been observed in animal reproduction studies.

3.Dicloxacillin crosses the placenta (Depp 1970). Maternal use of penicillins has generally not resulted in an increased risk of birth defects.

授 乳 期: 1.藥品會分泌至乳汁中,應小心使用。

2. Penicillins are considered compatible with breastfeeding when used in usual recommended doses (WHO 2002).

3.Dicloxacillin has been recommended to treat mastitis in lactating women (Amir 2014).