

Felodipine

【OFELO】Felpin[®] Extended Release Tablets 5mg

ATC Code : C08CA02

中文名：菲可平持續釋放膜衣錠 «信東»

適應症：高血壓、心絞痛。

藥理分類：**Calcium Channel Blocker, Dihydropyridine.**

用法用量：**Administration:** orally, without food or with a light meal. Swallow tablet whole; tablet should not be divided, crushed, or chewed.

Indications and dosage regimen:

Hypertension:

- Initially, 5 mg once daily. Adjust dosage at not less than 2 weeks intervals to achieve BP control.
- Maintenance, 5-10 mg once daily
- Hepatic Impairment & Elderly ($\geq$ 65 yrs):
usual initial dose, 2.5 mg daily; MAX 10 mg/day.

Angina pectoris:

5 to 10 mg once daily; if initiated at 5 mg, increase dose to 10 mg once daily as tolerated after 2 to 4 weeks.

Safety and efficacy not established in children.

不良反應：四肢水腫、頭痛、潮紅等。

注意事項：1. 錠劑應於早上服用，配以開水整顆吞服，不可剝半或磨碎服用。可於空腹時或輕膳（指不含高脂肪或高碳水化合物）後服用。
2. 本品應避免以葡萄柚汁送服。

禁忌：1. 懷孕
2. 已知對本品成分(Felodipine)或賦形劑過敏
3. 未代償性心衰竭
4. 急性心肌梗塞
5. 不穩定性心絞痛

交互作用：1. 增加 Felodipine 血中濃度：Cimetidine、Erythromycin、Itraconazole、Ketoconazole、葡萄柚汁。
2. 降低 Felodipine 血中濃度：Phenytoin、Carbamazepine、Rifampicin、Barbiturates、St. John's wort (聖約翰草)。

懷孕期：1. 懷孕期間不應給予 felodipine。
2. Calcium channel blockers may be used to treat hypertension in pregnant women; however, agents other than felodipine are more commonly used (ACOG 203 2019; ESC [Regitz-Zagrosek 2018]).

授乳期：1. 在乳汁中可被檢測出；授乳婦服用治療劑量，似乎不會影響嬰兒。
2. Due to the potential for serious adverse reactions in the breastfeeding infant, the manufacturer recommends a decision be made whether to discontinue breastfeeding or to discontinue the drug, considering the importance of treatment to the mother.