

Indomethacin

【OINDO】Indomen[®] 25mg/Cap

ATC Code : M01AB01

中文名：克炎膠囊 «永信»

適應症：關節炎、痺麻質斯(rheumatism)、變形性關節症等炎症之消炎、鎮痛、解熱以及其他疾患之消炎。

藥理分類：**Analgesic, Nonopioid; Nonsteroidal Anti-inflammatory Drug (NSAID).**

用法用量：**Administration:** orally immediately after meals or with food or antacids (to reduce adverse GI effects).

Indications and dosage regimens:

Inflammatory Diseases:

— Juvenile Rheumatoid Arthritis

Children 2 years of age:

Initially, 2 mg/kg daily in divided doses. Increase dosage until a satisfactory response is achieved, up to maximum dosage of 4 mg/kg daily or 150-200 mg daily (whichever is less) in divided doses. As symptoms subside, reduce dosage to the lowest effective level or discontinue the drug.

— Osteoarthritis, Rheumatoid Arthritis, or Ankylosing

Spondylitis:

Initially, 25 mg 2 or 3 times daily. If needed, increase dosage by 25 or 50 mg daily at weekly intervals until a satisfactory response is obtained up to a maximum dosage of 150-200 mg daily.

— Gout: 50 mg 3 times daily until pain is tolerable; then reduce dosage rapidly and discontinue.

— Painful Shoulder:

70-150 mg daily in 3 or 4 divided doses.

Discontinue once symptoms have been controlled for several days; usual course of therapy is 7-14 days.

Pericarditis:

Reduction of pain, fever, and inflammation of pericarditis; however, other drugs (i.e., aspirin) generally are preferred.

— Pediatric: 50-100 mg daily in 2-4 divided doses.

— Adults: 75-200 mg daily in 3 or 4 divided doses.

不良反應：噁心、消化不良、頭痛、眩暈。

注意事項：1.可與食物或牛奶一起服用。

2.如用藥後發生喉痛、口腔/黏膜潰爛、皮疹等症狀,宜立即就醫並考慮停藥。

懷 孕 期：懷孕後期應避免使用（因可能引起主動脈導管的早期癒合）。

授 乳 期：本品會分泌於乳汁中，不建議使用。