Phenazopyridine HCl

[OPYR] Pyridium (Urogen®) 50 mg/Tab ATC Code : G04BX06

中文名: 抑炎源膜衣錠 《強生》

適應症: 膀胱炎、腎盂炎、尿道炎、泌尿生殖器粘膜之鎮痛。

藥理分類: Analgesic, Urinary.

作用機轉: An azo dye which exerts local anesthetic or analgesic action on urinary tract mucosa

through an unknown mechanism.

用法用量: **Administration:** administer without regard to meals.

Indications and dosage regimen:

Dysuria, for symptomatic relief of pain, burning, urgency, and frequency arising from irritation of the urinary tract (infection, trauma, surgery, endoscopic procedures, or catheter placement):

-Adults:

2 Tablets TID.

-Children 6-12 yrs:

1 Tablet Tid.

不良反應: 噁心、腹痛、消化不良;頭暈、頭痛、發燒;皮膚或眼睛變黃;皮膚起疹; 臉、手指、腳腫脹等。

交互作用: ● May ↑ the adverse/toxic effect of **Sodium Nitrite**. Combinations of these agents may increase the likelihood of significant methemoglobinemia.

• Nitric Oxide may ↑ the adverse/toxic effect of Methemoglobinemia Associated Agents. Combinations of these agents may increase the likelihood of significant methemoglobinemia.

注意事項: 1、本品不可研磨使用。(避免沉積牙齒導致著色)

2、服用本劑後,尿液會呈黃色或黃褐色,屬正常現象,不必擔心。

懷孕期: Phenazopyridine 會通過人類胎盤。動物試驗未發現不良作用。

授 乳 期: 本品用於授乳婦時,其對嬰兒的安全性尚未確立。