Sucralfate hydrate

[OULSG] Ulban A Gran® 900mg/g, 1g/Sachet ATC Code: A02BX02

中文名: 胃樂伴顆粒 《Towa》

適應症: 胃潰瘍及十二指腸潰瘍。

Gastrointestinal Agent, Miscellaneous. 藥理分類:

作用機轉: Forms a complex by binding with positively charged proteins in exudates, forming a

viscous paste-like, adhesive substance. This selectively forms a protective coating that acts locally to protect the gastric lining against peptic acid, pepsin, and bile salts.

用法用量: **Administration:** orally on an empty stomach 1 hour before meals and at bedtime.

Indications and dosage regimens:

Duodenal ulcers:

- Treatment:

1 g 3-4 times daily OR 2 g BID for 4-8 weeks.

-Maintenance: 1 g BID

Gastric ulcers, maintenance: 1 g BID.

GI haemorrhage:

-Treatment:

1 g 6 times/day. Max 8 g /day.

便秘、口渴、噁心、嘔氣。 不良反應:

注意事項: 1. 空腹(飯前1小時) 或睡前服用。

> 2. 對腎障礙患者(長期投藥時可能出現 Dialysis Encephalopathy 和 Dialysis Osteomalacia, 需定期檢查血中的鋁、磷、alkaliphosphatase 等)。

3. 缺乏磷酸鹽的患者(鋁在消化道內和磷酸鹽結合,阻礙其吸收)謹慎投與。

交互作用: 1.本劑以空腹服用為宜,如必須與制酸劑併服,兩者須至少錯開半小時以上。

2.本劑會阻礙 Quinolone 類抗菌劑的吸收,兩者不得併服。

3.本劑會阻礙 tetracycline 類抗菌劑的吸收,兩者不得併服。

- **Baloxavir** Marboxil: ↓ the serum concentration of Baloxavir Marboxil.
- **Antacids**: ↓ the therapeutic effect of Sucralfate.
- **Bisphosphonate**(oral): ↓ the serum concentration of Bisphosphonates.
- **Digoxin**: ↓ the serum concentration of Digoxin.
- Furosemide (oral): ↓ the serum concentration of Furosemide.

 Quinolones (oral): ↓ the serum concentration of Quinolones.

 Sulpiride: ↓ the serum concentration of Sulpiride.

- **Tetracyclines**: ↓ he absorption of Tetracyclines.

1. Sucralfate is only minimally absorbed following oral administration. Based on 懷 孕 期: available data, sucralfate does not appear to increase the risk of adverse fetal events when used during the first trimester.

> 2. Sucralfate may be used for the treatment of duodenal ulcer or reflux in pregnancy (Cappell 2003; Dağlı 2017; Richter 2007).

1.It is not known if sucralfate is present in breast milk. 授乳期:

- 2. Sucralfate is only minimally absorbed following oral administration.
- 3. Although the manufacturer recommends that caution be exercised when administering sucralfate to breastfeeding women, use is considered acceptable.