Ursodeoxycholic Acid

[OURSU] Urso (Urose®) 300mg/Cap

中文名: 幽立平膠囊 《意欣》

適應症: 膽固醇系膽結石之溶解、原發性膽道肝硬化 (primary biliary cirrhosis, PBC) 之

ATC Code: A05AA02

肝功能改善。

藥理分類: Gallstone Dissolution Agent.

用法用量: Administration: should be taken with meals or a snack since ursodiol dissolves more

rapidly when bile and pancreatic juice are present in the intestinal chyme.

Indications and dosage regimen:

**Disease of liver, Cholestatic:** 

13-15 mg/kg/day, divided into 2 doses (morning and bedtime).

**Gallstone:** 

8-10 mg/kg/day, divided into 2-3 doses.

Gallstone, During rapid weight loss; Prophylaxis:

300 mg 2 times a day (morning and bedtime)

**Hepatitis, chronic:** 

450-600 mg/day in divided doses.

Primary biliary cirrhosis:

13-15 mg/kg/day (in 4 divided doses)

Congenital dilatation of lobar intrahepatic bile duct:

15-20 mg/kg/day in divided doses

Safety and effectiveness not established in children.

不良反應: 腹瀉、便秘、消化不良、頭痛、金屬味覺、脹氣、背痛等。

注意事項: 1.猛爆性肝炎、膽道閉塞患者禁用本劑。

2.飯中或飯後服用。本品不可與含鋁的制酸劑並用。

3.本藥藥效產生慢,可能需要幾個月的時間才能將膽結石完全溶解。接受本藥 治療期間應定期檢查肝功能。

4. 與經口糖尿病藥(Tobulamide)等併用,作用增強。

懷 孕 期: 1.妊娠婦女不要使用。(仿單)

2. Adverse events have not been observed in animal reproduction studies. Ursodiol (ursodeoxycholic acid) is the treatment of choice for intrahepatic cholestasis of

pregnancy (Kremer 2011).

授 乳 期: It is not known if ursodiol is excreted in breast milk. The manufacturer recommends that caution be exercised when administering ursodiol to breastfeeding women.