Orlistat

【OXEN】Xenical® 120mg/Cap *ATC Code：A08AB01*

中 文 名： 羅鮮子膠囊 «Roche»

適 應 症： 低卡洛里飲食，適合肥胖病患的治療，包括有與肥胖相 關危險因子之病患。

藥理分類： Lipase Inhibitor

用法用量： **Administration:**

- Orally, administer **during or up to 1 hour after** each main meal containing fat.
- Separate dose by **at least 2 hours** from multivitamin containing fat-soluble vitamins.
- Omit dose if a meal is missed or contains no fat.

Dosage regimens:

120 mg 3 times daily with each main meal containing fat.

No additional benefit with dosages > 120 mg 3 times daily.

Safety and efficacy not established in children < 18 yr.

不良反應： 油斑（沾褲）、胃脹氣、急便、油便、排油、排便增加、腹痛、脹氣。

交互作用：

1. **Anticonvulsants**: ↓ anticonvulsant effectiveness.
2. Ritonavir (**Paxlovid**®含此成分): may result in loss of virological control.
3. **Anticoagulants**: ↑ risk of bleeding.

注意事項： 1.治療期間不得超過 2 年。

2.若不進食或此餐不含脂肪時則可省略服用此藥。

3.可能影響脂溶性維生素 A、D、E、K 的吸收。兩者服用時間需相隔至少 2 小時。

4.若某一餐含脂量特別高時服用，則胃腸道的副作用會增加。

5.建議主餐進行中或最遲進餐後 1 小時內服用。

懷 孕 期： 動物實驗中並未發現 orlistat 有胚胎毒性或致畸胎效應。在無動物之致畸胎效應下，於人類中亦不預期有致畸胎的反應。然而，在缺少臨床數據下，並**不建議**懷孕 時服用 Xenical®。

授 乳 期： 尚無 orlistat 是否會分泌至人類乳汁的研究。哺乳時**不應服用** Xenical®。

備 註： 本品健保不給付，自費處方計價單位量為 42 粒。