Levocetirizine 2HCl

OXYI Xyzine[®] F.C. Tablets 5mg/Tab

中文名: 萊欣膜衣錠5毫克 《中國化學》

適應症: 治療成人及六歲以上孩童因過敏性鼻炎、慢性蕁麻疹等所引起的各種過敏徵狀。

藥理分類: Histamine H1 Antagonist, Second Generation; Piperazine Derivative.

用法用量: Administration: orally, swallowed with liquid and may be taken with or without

 $ATC\ Code : R06AE09$

food. It is recommended to take the daily dose in one single intake.

Dosage regimens:

Adults & Children ≥ 6 yr:

5 mg once daily.

Dosage adjustment in Patients with renal impairment:

Group	Clcr (mL/min)	Dosage & Frequency
Normal	80	5mg once daily
Mild	50-79	5mg once daily
Moderate	30-49	5mg once every 2 days
Severe	< 30	5mg once every 3 days
End-stage renal disease- Patients undergoing dialysis	<10	Contra-indicated

禁 忌: 1.對 piperazine 衍生物過敏者,應避免使用本劑。

- 2.嚴重腎功能障礙者(Clcr < 10mL/min)禁用。
- 3.若具有對 galactose 無耐受力,或 Lapp lactase 缺乏,或有 glucose-galactose 吸收障礙等遺傳疾病,則禁用本劑。

不良反應: 嗜睡(5.2%)、頭痛(2.6%)、口乾(2.6%)、疲勞(2.5%)等。

交互作用:

- Azelastine (Nasal): † the CNS depressant effect of Levocetirizine.
- Levocetirizine may \(\bar{} \) the anticholinergic effect of **Glycopyrrolate** (Oral Inhalation), **Tiotropium**.
- **Ipratropium** (Oral Inhalation), **Umeclidinium, Amantadine**: ↑ the anticholinergic effect of Levocetirizine.
- Levocetirizine may \(\) the ulcerogenic effect of **Potassium Chloride**.
- Acetylcholinesterase Inhibitors (Donepezil, Edrophonium, Galantamine, Neostigmine, Physostigmine, Pyridostigmine, Rivastigmine): ↓ the therapeutic effect of Levocetirizine. Levocetirizine: ↓ the therapeutic effect of Acetylcholinesterase Inhibitors.

注意事項: 1.6歲以下孩童不建議使用本劑。

- 2.飯前或飯後服用均可。
- 3.勿併服含酒精飲 料。
- 4.服用此藥後,勿駕車或操作危險器械。

懷 孕 期: 1.Second generation antihistamines may be used for the treatment of allergic rhinitis and urticaria during pregnancy;

2.Information related to the use of levocetirizine in pregnancy is limited and other medications may be preferred (Wallace 2008; Zuberbier 2018).

授 乳 期: 1. It is not known if levocetirizine is present in breast milk. Levocetirizine is the active enantiomer of cetirizine which is present in breast milk

2. When treatment with an antihistamine is needed in breastfeeding women, second generation antihistamines are preferred (Butler 2014; Powell 2015; Zuberbier 2018).

過量處置: 1.若服用過量時,成人會有嗜睡現象,孩童則會先興奮不安而後伴隨睏倦。

2.對於 levocetirizine 過量並無特殊的解毒劑,建議以徵狀治療,亦可考慮洗胃 (服用後短期內)。血液透析對 levocetirizine 的排除無效。