Acyclovir

【TZOV】眼用 Devirus® Oph. Ointment 3% 4.5g/Tube ATC Code: S01AD03

中文名: 力克眼用軟膏 《溫士頓》

適應症: 單純疱疹性角膜炎。

藥理分類: Antiviral Agent, Ophthalmic.

用法用量: Administration:

For topical ophthalmic use only; avoid touching tip of applicator to eye(s) or other surfaces. After application close the eye(s) for 1 to 2 minutes; excess ointment may be wiped away.

Dosage regimen:

The dosage for all age groups is the same.

A 10 mm ribbon of the ointment should be placed inside the lower conjunctival sac 5 times a day at 4 hours intervals; continue for at least 3 days after healing.

不良反應: 可能會引起眼睛短暫性的輕微刺痛、癢、灼熱感。

注意事項: 打開一個月後之本品必須丟棄。

懷 孕 期: 1.動物試驗未發現胎兒毒性或畸形。尚無足夠之人體安全資料。

2. 唯有效益高於可能風險時才使用。

授 乳 期: 1.目前有限的數據顯示,口服本藥會由乳汁分泌出來,眼用則未知。

2. therapy should consider the risk of infant exposure, the benefits of breastfeeding to

the infant, and benefits of treatment to the mother.

3. Systemic use of acyclovir is considered compatible with breastfeeding (WHO

2002).

储 存: 25 ℃以下储存。經開瓶一個月後之本品必須丟棄。